

Connecting to Jaime - Dialog - 245985

Subaccount = 031749/268956-JG_071207

[File 351] **Derwent WPI** 1963-2007/UD=200743

(c) 2007 The Thomson Corporation. All rights reserved.

[File 345] **Inpadoc/Fam.& Legal Stat** 1968-2007/UD=200727

(c) 2007 EPO. All rights reserved.

**File 345: Preview the enhanced INPADOC database in ONTAP File 253. For more information, visit www.dialog.com/inpadoc.*

[File 331] **Derwent WPI First View UD=200743** (c) 2007 The Thomson Corp.

. All rights reserved.

**File 331: For patent family information, search also File 351, 352, or 350.*

[File 347] **JAPIO** Dec 1976-2007/Dec(Updated 070702)

(c) 2007 JPO & JAPIO. All rights reserved.

S1 3 S PN=JP 6219967

1/3,LS,AB/1 (Item 1 from file: 351)
Derwent WPI
(c) 2007 The Thomson Corporation. All rights reserved.

0006916623

WPI Acc no: 1994-312661/

XRAM Acc no: C1994-141928

New peptide-lipid derivs. bound directly or via linker to lipid - useful for inhibiting mouse lung cancer cell line, 3LL cell adhesion by fibronectin

Patent Assignee: DDS KENKYUSHO KK (DDSK-N)

Inventor: ISHIKURA T; KANEKO H; KATO T; MURAHASHI N; NAGASO H; SASAKI A; TANAKA I

Patent Family: 2 patents, 1 countries

Patent Number	Kind	Date	Application Number	Kind	Date	Update	Type
JP 6219967	A	19940809	JP 19939290	A	19930122	199439	B
JP 2579730	B2	19970212	JP 19939290	A	19930122	199711	E

Priority Applications (no., kind, date): JP 19939290 A 19930122

Patent Details

Patent Number	Kind	Lan	Pgs	Draw	Filing Notes		
JP 6219967	A	JA	12	1			
JP 2579730	B2	JA	12	0	Previously issued patent	JP 06219967	

Alerting Abstract JP A

Peptides contg. an amino acid sequence, Arg-Gly-Asp (RGD sequence) bound directly or via linker to lipid are new.

The exogenous peptides contg. RGD peptide can bind to cancer cells, competing with fibronectin, and inhibit intercellular adhesion by fibronectin. RGD sequence has been observed to suppress transfer of cancer cells (Humphries M.J., et al., Science, 2, 33, 467 (1986)).

USE - The liposome contg. RGD-lipid derivatives can be used for effectively inhibiting mouse lung cancer cell line, 3LL cell adhesion by fibronectin.

Basic Derwent Week: **199439**

1/3,LS,AB/2 (Item 1 from file: 345)
Inpadoc/Fam.& Legal Stat
(c) 2007 EPO. All rights reserved.

Acc no: 11932139

Basic Patent (No,Kind,Date): JP 6219967 A2 19940809

No. of Patents: 002

PEPTIDE-LIPID DERIVATIVE AND LIPOSOME (English)

Patent Assignee: D D S KENKYUSHO KK

Author (Inventor): ISHIKURA TOYOAKI; SASAKI ATSUSHI; NAGASO HIROSHI;

MURAHASHI NAOICHI; KATO TAKASHI; KANEKO HIDEO; TANAKA ISAO

IPC: *A61K-047/48; A61K-009/127; C07K-015/12

Language of Document: Japanese

Patent Family:

Patent No	Kind	Date	Applic No	Kind	Date	
JP 6219967	A2	19940809	JP 939290	A	19930122	(BASIC)
JP 2579730	B2	19970212	JP 939290	A	19930122	

Priority Data (No,Kind,Date):

JP 939290 A 19930122

1/3,LS,AB/3 (Item 1 from file: 347)
JAPIO
(c) 2007 JPO & JAPIO. All rights reserved.
04548067 **PEPTIDE-LIPID DERIVATIVE AND LIPOSOME**

Pub. No.: 06-219967 [JP 6219967 A]
Published: August 09, 1994 (19940809)

Inventor: ISHIKURA TOYOAKI
SASAKI ATSUSHI
NAGASO HIROSHI
MURAHASHI NAOICHI
KATO TAKASHI
KANEKO HIDEO
TANAKA ISAO

Applicant: D D S KENKYUSHO KK [000000] (A Japanese Company or Corporation), JP (Japan)
Application No.: 05-009290 [JP 939290]
Filed: January 22, 1993 (19930122)
Journal: Section: C, Section No. 1271, Vol. 18, No. 588, Pg. 131, November 10, 1994
(19941110)

ABSTRACT

PURPOSE: To provide a new peptide-lipid derivative useful as a metastasis- suppressing substance for cancer cell.

CONSTITUTION: A lipid is bonded directly or via a linker to a peptide containing a sequence composed of Arg-Gly-Asp. The total polymerization degree of the amino acids in the peptide is ≤ 20 and the lipid to be used for the production of the peptide-lipid derivative is preferably cholesterol and 8-18C alkyl group. The bonding of the lipid to the peptide containing the Arg-Gly-Asp sequence can be carried out e.g. by reacting an amino group or a carboxyl group of a peptide with a cholesterol derivative or an alkyl group having a functional group capable of forming a covalent bond with the functional group of the peptide. A liposome can be prepared by compounding the peptide-lipid derivative.